

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/101825/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Patrício, AR, Formia, A, Barbosa, C, Broderick, AC, Bruford, Michael William ORCID: <https://orcid.org/0000-0001-6357-6080>, Carreras, C, Catry, P, Ciofi, C, Regalla, A and Godley, BJ 2017. Dispersal of green turtles from Africa's largest rookery assessed through genetic markers. Marine Ecology Progress Series 569 , pp. 215-225. 10.3354/meps12078 file

Publishers page: <http://dx.doi.org/10.3354/meps12078>
<<http://dx.doi.org/10.3354/meps12078>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.



Dispersal of green turtles from Africa's largest rookery assessed through genetic markers

Running page head: **Green turtle connectivity in the Atlantic**

Ana R. Patrício ^{1,2,*}, Angela Formia^{3,4}, Castro Barbosa ⁵, Annette C. Broderick ¹, Mike Bruford ⁶, Carlos Carreras ^{7,8}, Paulo Catry ², Claudio Ciofi ⁴, Aissa Regalla ⁵, Brendan J. Godley¹

¹ Centre for Ecology and Conservation, University of Exeter, TR10 9EZ, Penryn, UK

² MARE – Marine and Environmental Sciences Centre, ISPA – Instituto Universitário, 1140-041, Lisbon, Portugal

³ Wildlife Conservation Society, Marine Program, BP 7847, Libreville, Gabon

⁴ Department of Biology, University of Florence, Sesto Fiorentino, 50019, FI, Italy

⁵ Institute of Biodiversity and Protected Areas of Guinea-Bissau (IBAP), CP – 70, Bissau, Guinea Bissau

⁶ School of Biosciences, Cardiff University, CF10 3AX, Cardiff, UK

⁷ University of Barcelona, Department of Genetics, Microbiology and Statistics, 08028, Barcelona, Spain

⁸ Institute of Biodiversity Research of Barcelona, IRBio, 08028, Barcelona, Spain

Abstract

Marine turtles are highly migratory species that establish multiple connections among distant areas, through oceanic migration corridors. To improve the knowledge on the connectivity of Atlantic green turtles, we analysed the genetic composition and contribution to juvenile aggregations of one of the world's largest rookeries at Poilão Island, Guinea-Bissau. We amplified 856bp mitochondrial DNA (mtDNA) control region sequences of this population (n=171) containing the ~490bp haplotypes used in previous studies. Haplotype CM-A8 was dominant (99.4%) but it divided in two variants when the whole 856 bp was considered: CM-A8.1 (98.8%) and CM-A8.3 (0.6%). We further identified the haplotype CM-A42.1 (0.6%), found previously only in juvenile foraging grounds at Argentina, Brazil and Equatorial Guinea. The Poilão breeding population was genetically different from all others in the Atlantic (F_{ST} range 0.016-0.961, $P < 0.001$). An extensive 'Many-to-many' mixed-stock analysis (MSA) including 14 nesting populations (1,815 samples) and 17 foraging grounds (1,686 samples) supported a strong contribution of Poilão to West Africa (51%) but also to Southwest Atlantic (36%). These findings, in particular the strong connectivity within West Africa, where illegal harvesting is still common, should motivate conservation partnerships, so that population protection can be effectively extended through all life-stages. Our study expands the knowledge on migration patterns and connectivity of green turtles in the Atlantic, evidences the importance of larger sample sizes and emphasises the need to include more finely resolved markers in MSAs and more genetic sampling from West African foraging grounds to further resolve the connectivity puzzle for this species.

Key-words: connectivity, dispersal, green turtle, migration, mitochondrial DNA (mtDNA), mixed-stock analysis (MSA), population genetics, West Africa

* R.Patricio@exeter.ac.uk

Introduction

Many marine species undertake migratory movements among distant geographic areas and across distinct habitats, for feeding, reproduction or development. As a result they may be subject to a diverse range of threats during their extensive movements. Sea birds (Catry et al., 2011), marine mammals (Rasmussen et al., 2007), large fish (Bonfil et al., 2005, Rooker et al., 2014) and sea turtles (Hays and Scott, 2013) undertake such movements and are known to play important ecological roles. Understanding their dispersal patterns and the links they establish among different areas is critical to contextualize threats and inform effective management strategies (Rees et al., 2016).

Marine turtles are long-lived organisms and their life histories are marked by ontogenic habitat shifts and large-scale migrations (Bowen & Karl, 2007). Green turtles (*Chelonia mydas* L) associate with oceanic currents after hatching and undergo an oceanic pelagic stage, which is thought to last ca. 3 – 5 years (Reich et al., 2007). After this period, often referred to as ‘the lost years’, as the whereabouts of the turtles at this phase are poorly known, they generally recruit to coastal habitats, which may change seasonally (Fukuoka et al., 2015), and shift into benthic foraging at a straight-carapace-length of 25 – 35 cm (Bolten, 2003). These neritic zones are used as developmental habitats and turtles may spend several years foraging in the same area until reaching a size or maturity stage that triggers them to migrate to additional foraging areas (Patrício et al., 2011, Patrício et al., 2014, Shimada et al., 2015). Upon reaching maturity, adults make periodic migrations between their neritic foraging areas and natal rookeries (Bowen and Karl, 2007). This complex migratory behaviour creates multiple connections among distant coastal areas through oceanic migration corridors (Velez-Zuazo et al., 2008). Genetic studies have been critical in enlightening such connectivity (Encalada et al., 1996; Naro-Maciel et al., 2007; Prosdocimi et al., 2012).

Most studies have used sequences of the control region of the mitochondrial DNA (mtDNA), a maternally inherited genetic marker (Bowen and Karl, 2007). This marker shows generally high levels of genetic structuring among marine turtle nesting populations worldwide, supporting the natal homing hypothesis, in which the females of marine turtles return to the beaches where they were born to reproduce as a consequence of philopatry (Meylan et al., 1990). In contrast, foraging aggregations are usually mixed stocks composed of individuals from different rookeries (Bowen and Karl, 2007). The high genetic structuring of nesting populations allows the use of mixed stock analysis (MSA, Millar, 1987), to estimate contributions of rookeries (stocks) to mixed foraging grounds (mixed stocks). A Bayesian MSA (Pella and Masuda, 2001) has been widely applied, allowing the incorporation of informative priors, such as rookery size or geographic distance. Bolker et al (2007) subsequently developed a 'many-to-many' mixed stock analysis (m2m MSA), aiming to simultaneously answer the questions 1) where do the individuals from a given source population go? and 2) where do individuals from a given mixed foraging ground originate? Limitations of MSAs have been pointed out however, in particular the assumption that all source populations and mixed aggregations have been adequately sampled (Proietti et al., 2012). The existence of orphan haplotypes at juvenile foraging grounds indicates that some stocks still lack genetic assessment or have not yet been adequately sampled; hence estimates should be interpreted cautiously and along with meaningful ecological data.

One controversial result of recent MSAs of the Atlantic green turtles is the suggested potential connectivity between Guinea-Bissau, West Africa, and the Southwest Atlantic. Although MSAs have supported this migration (Bolker et al., 2007; Monzón-Argüello et al., 2010; Naro-Maciel et al., 2012), the fact that the population at Poilão, Guinea-Bissau, was found to be fixed for the common South Atlantic haplotype (CMA-8, Encalada et al., 1996; Formia et al., 2006; Godley et al., 2010) has limited the interpretations of these results. Notably, the discovery of exclusive haplotypes at low frequency is highly dependent on sample size. This putative migration seems to involve movements greater than expected,

according to the 'closest to home' hypothesis where immature turtles tend to move to and settle in foraging grounds closest to their natal beach after recruiting to neritic habitats (Bolker et al., 2007). Additionally, studies using particle dispersal modelling with major oceanic currents did not support this connectivity (Godley et al., 2010; Putman and Naro-Maciel 2013). However, when Putman and Naro-Maciel (2013) estimated the origins of the green turtle Atlantic mixed stocks, tracking particles back through time, this crossing seemed feasible, albeit at low incidence. Lagrangian drifter data have further shown this route to be possible with particle drift (Monzón-Argüello et al., 2010; Proietti et al., 2012). Finally, a similarly large-scale migration of post-hatchling green turtles from Suriname to Cape Verde was supported using mtDNA (Monzón-Argüello et al., 2010).

With this in mind we investigate two questions: 1) where do the post-hatchlings from Poilão disperse to?, and 2) do some of the juveniles found at Southwest Atlantic foraging grounds originate in Poilão? To answer these questions we greatly increased the available sample to characterize the genetic composition of Poilão's nesting population, in an attempt to detect rare haplotypes. We then sought to improve our understanding of the migration patterns and connectivity among Atlantic green turtle populations by comparing our results with molecular data (n = 3,501 sequences) from 14 nesting populations and 17 foraging grounds, resulting in the most extensive analysis thus far for this species in the Atlantic.

Methods

Study site and sampling

Poilão Island (N10°52', W15°43') is part of the João Vieira and Poilão Marine National Park (PNMJVP), in the Bijagós Archipelago, Guinea-Bissau. It hosts one of the major green turtle nesting populations worldwide (Catry et al., 2002, 2009). This population has been monitored yearly around the peak of the nesting season (August -September) since 2004. In 2013 and 2014 we collected skin samples from 171 nesting females. Samples were taken from the shoulder area using a 6 mm sterile biopsy punch as the females laid their eggs and stored in

96 % ethanol at room temperature. All sampled individuals were identified with unique tags on both front flippers to avoid sample duplication. Furthermore, the loss of a metal tag leaves scar marks easily recognized within, so we were certain that no previously tagged individual was mistakenly identified as 'new'. Sampling protocols were approved by the research ethics committee of the University of Exeter and the government of the Republic of Guinea-Bissau.

Sequencing and haplotype assignment

We extracted DNA using the QIAGEN® DNeasy blood & tissue kit, according to the manufacturer's instructions. A fragment of ~860bp of the mtDNA control region was amplified in a polymerase chain reaction (PCR) with the primers LCM15382 (5'-GCTTAACCCTAAAGCATTGG-3') and H950 (5'-TCTCGGATTTAGGGGTTT-3') (Abreu-Grobois et al. 2006) which includes the short region (~486 bp) historically surveyed for green turtle genetic studies (Bjorndal et al., 2006; Encalada et al., 1996; Formia et al., 2007; Lahanas et al., 1998). Amplifications were performed in a total volume of 25 µl, containing 2.5 µl of Taq buffer, 3 µl of dNTPs, 1 µl of MgCl₂, 0.5 µM of each primer at 10 µM, and 0.2 µl of Taq DNA polymerase. Cycling conditions were 94° C for 5 min, followed 35 cycles at 94° C for 1 min, 55° C for 1 min and 72° C for 1 min with a final extension step at 72° C for 10 min. Desired PCR products were purified with a combined Exonuclease I and Shrimp Alkaline Phosphatase solution (ExoSAP®). The reaction was incubated for 15 min at 37°C, followed by 15 min incubation at 80°C to inactivate the two enzymes. Sequences of forward and reverse DNA strands were performed at MacroGen Inc. (Netherlands). Sequences were assembled and aligned manually using BioEdit 7.2.5 (Hall 1999). Unique haplotypes were identified using the Basic Local Alignment Search tool (BLAST) from the National Centre for Biotechnology information (<http://www.ncbi.nlm.nih.gov/>), following the nomenclature of the Archie Carr Center for Sea Turtle Research, ACCSTR; <http://accstr.ufl.edu/ccmtDNA.html>.

Population structure

To assess the genetic diversity of the nesting population at Poilão compared with the other Atlantic nesting populations we truncated the mtDNA fragments to 490 bp length, the fragment historically explored and for which most genetic information of other locations is currently available. We used Arlequin 3.5.1.3 (Excoffier and Lischer 2010) to estimate the haplotype (h) and nucleotide (π) diversity of nesting populations, to estimate the genetic distances among population pairs (Φ_{st}) and to test the significance of differentiations with exact tests based on haplotype frequencies. A false discovery rate (FDR) correction (Narum, 2006) was applied to calculate the most fitting threshold for the P-value significance considering the number of comparisons involved in the analysis and under an expected original threshold of $P < 0.05$. To contextualize our sampling location within the Atlantic region, the genetic distances were used to perform a principal coordinate analysis (PCoA) using the package GenALEX 6.5.0.1 (Peakall and Smouse, 2012). We tested the significance of the PCoA grouping with an AMOVA, using Arlequin 3.5.1.3 (Excoffier and Lischer 2010).

‘Many-to-many’ Mixed-stock analysis

We generated a dataset of 14 nesting populations ($n=1,815$) and 17 foraging grounds ($n=1,686$) when including our new mtDNA data for Poilão to the previously existing data for Atlantic nesting populations and foraging grounds (see figure 1 for sites included in this study and literature sources). We used only sequences generated by this study to characterize the genetic composition of Poilão in order to avoid potential pseudoreplication with datasets obtained in previous years. Relative contributions to foraging areas from nesting populations (mixed stock-centric approach), and probable use of foraging grounds from nesting populations (source-centric approach) were estimated with m2m MSA, using the R package mixstock (Bolker et al., 2007) and WinBUGS (Lunn et al., 2000). We conducted the MSA including the number of nesting females in each population (Seminoff et al., 2015) as a weighting factor (Prosdocimi et al., 2012). We used the Gelman-Rubin diagnostic to assess convergence of the chains to the posterior distribution, assuming that

there was no evidence of non-convergence at values < 1.2 (Pella and Masuda, 2001). As it is reasonable to assume that other African juvenile aggregations remain to be identified we simulated a juvenile foraging ground fixed for haplotype CM-A8 (similar to Naro-Maciel et al 2012), with a sample size equal to the mean of the foraging grounds sample sizes ($n=99$), and added this sample to the dataset to conduct another m2m MSA, as described above.

Results

Genetic composition of Poilão

Genetic variability of the Poilão nesting population was the lowest of all Atlantic populations ($h \pm SD = 0.012 \pm 0.011$, $\pi \pm SD = 0.0001 \pm 0.0003$, table 1). The haplotype CM-A8 was dominant as suggested by previous studies (Formia et al., 2006). However the use of longer sequences (856bp sequences) distinguished two variants of this haplotype: CM-A8.1 (98.8 %) and CM-A8.3 (0.6 %). We also identified the haplotype CM-A42.1 (0.6 %), a previously orphan haplotype found to date only in juveniles from West Africa and South American foraging aggregations (see Table S1 for haplotype frequencies of nesting populations). Because this is a rare haplotype and not previously detected in the population we performed two independent PCRs, and sequenced the amplified fragment in two independent occasions, to confirm that this result was not a product of genotyping error.

Population Structure

The nesting population at Poilão was significantly different from all other Atlantic green turtle rookeries (Table S2). All other nesting populations were distinct from each other except when comparing Ascension Island with Bioko Island, Aves with Suriname, and Aves with Buck Island. The comparisons between Suriname and Buck Island, and between Sao Tome and Principe and Bioko became non-significant after FDR correction. Populations pairs where genetic differentiation was not detected were kept as discrete sources for the m2m MSA, based on their divergence in population size and geographic position (Monzón-Argüello et al., 2010; Putman and Naro-maciel, 2013). The PCoA separated rookeries by

region and evidenced three major groups: South Atlantic, Southeast Caribbean and Northwest Caribbean (Fig. 2), each group defined by a major haplotype(s): CM-A8, CM-A5 and CM-A3/A1, respectively. An accumulated 85.5% of the genetic variability was explained by the two principal coordinates of the PCoA. Although located in the North Atlantic, Poilão clustered within the South Atlantic group. Using this *a-priori* grouping in the AMOVA, highly significant structure was observed among the three groups ($F_{ST} = 0.691$, $P < 0.001$), with 55.9% of the variation found among groups.

‘Many-to-many’ Mixed-stock analysis

The source-centric m2m MSA indicated that most of Poilão’s hatchlings recruit to African foraging grounds (51.4%), but 36.2% would reach juvenile aggregations in the Southwest Atlantic and 8.6% reached North Atlantic aggregations (Fig. 3). A small proportion of the Poilão rookery was attributed to an ‘unknown’ foraging area (3.7%). The foraging ground-centric m2m MSA estimated that at Sao Tome, Corisco Bay and ‘West Africa’ (Liberia to Benin) foraging grounds, over 60% of the juveniles originate at Poilão, as do 31% of the green turtles foraging at Cape Verde (Fig. 4). Notably, at the Southwest Atlantic foraging aggregations proportions ranging from 16 – 41% were attributed to Poilão (Fig. 4). Adding the simulated West African foraging ground did not change contributions at a regional scale, but the relative contributions to the Gulf of Guinea were significantly lower (8 to 14 % lower, Fig. S1), to accommodate a large contribution to this putative aggregation. Because CM-A42 is a rare haplotype and therefore difficult to detect when sampling a population, we decided to run two additional MSAs using simulated datasets, each of these including haplotype CM-A42 in one of the other two major green turtle rookeries in the Atlantic (i.e. Costa Rica and Ascension Island), and observed no significant changes (Fig. S1).

Discussion

One of the principal techniques that can offer insight into the migratory connectivity of species with complex life cycles is genetics. The robustness of subsequent inferences,

however, are highly dependent on the amount of information available, including the number of populations and foraging grounds analysed, and the strength of the signal, including sample sizes at each site and length of the genetic sequence and number of genetic markers analysed. Here we substantially increased the sampling effort at one of the largest Atlantic green turtle rookeries, in Poilão, Guinea-Bissau, in order to resolve the uncertainties surrounding the connectivity between this nesting population and distant juvenile aggregations. We successfully found the origin of a previously orphan haplotype, present in West Africa but also in South American foraging grounds, giving strength to the hypothesis of east-to-west connectivity.

Post-hatchling dispersal to east and west

The contributions estimated by the m2m MSA confirm a strong connectivity within West Africa, as previously hypothesized (Godley et al., 2010), particularly with foraging grounds in the Gulf of Guinea (i.e. 'Sao Tome', 'West Africa' and 'Corisco'). This dispersal was also predicted under an ocean circulation model and through passive drifting associated with the Guinea current (Putman and Naro-Maciel, 2013). Due to the large size of the nesting population at Poilão, it is likely however that significant proportions of other African juvenile aggregations originate there. In Guinea-Bissau there are at least two known aggregations of immature green turtles; i) at Unhocomo and Unhocomozinho Islands, in the Bijagós Archipelago, ca. 100 km NE from Poilão Island, and ii) at Varela beach, ca. 200 km NE from Poilão, that have not been genetically described. The same is true for a foraging ground in Mauritania, mentioned in Godley et al (2010), and in Congo. We have shown that the estimated proportions of post-hatchlings distributed among West African foraging grounds depend on the inclusion of new juvenile aggregations. To fully understand the connectivity of the large nesting population at Poilão it is essential that investigation into identifying and genetically characterizing these aggregations is undertaken. The MSA also suggests the existence of a transatlantic developmental migration for the green turtle, from east to west,

potentially associated with the Equatorial currents, and continuing south, reaching foraging grounds in the south of Brazil and in Argentina.

Studies using estimations of passive drift with major oceanic currents to predict the movements of post-hatchlings have suggested that dispersal from Guinea-Bissau to Southwest Atlantic is unlikely (Godley et al., 2010; Putman and Naro-Maciel, 2013). However, marine turtle hatchlings are capable of oriented swimming significantly impacting trajectories (Putman et al., 2012a, 2012b; Scott et al., 2012), and able to swim against currents (Booth, 2014). Indeed, recent research has shown that drifter tracks can diverge substantially from those of young turtles (Putman and Mansfield, 2015) and it is likely that this process is contributing to observed divergence between genetic- and drift-based predictions (Naro-Maciel et al., 2016). Because CM-A42 is a rare haplotype and therefore difficult to detect, we ran additional MSAs using simulated datasets, including this haplotype in each of the two other major green turtle rookeries in the Atlantic (i.e. Costa Rica and Ascension Island), and observed no significant changes (Fig. S1).

Expanded sample size and geographic coverage

Formia et al (2006) assessed the genetic composition of Poilão nesting females (n=51) and found it was fixed for the South Atlantic dominant mtDNA haplotype CM-A8. By extending this previous sample size, we were able to detect a rare haplotype, CM-A42, which to date had only been reported from juvenile green turtles foraging in South America, and in West Africa. This enabled the differentiation of Poilão from other Atlantic rookeries, agreeing with the high philopatry, characteristic of the green turtle, and the fine scale differentiation existent in other places. Increasing sample size has previously been shown to improve statistical power of detection of structure among populations, through the finding of rare haplotypes (Formia et al., 2007).

The existence of non-significant comparisons among certain population pairs could result from i) recent isolation, such that haplotype frequencies did not have time to differentiate, or ii) current gene flow, mediated by incidental deviations from natal homing. Lack of differentiation between Bioko and Ascension Island has been attributed to recent colonization of the former (Formia et al., 2006). Likewise, Aves and Buck Island may be more recent than the more diverse population in Suriname. Alternatively, the proximity between Aves and Buck Island (<300 km), and between Bioko and Sao Tome (<400 km), may be more likely to result in occasional migrants preventing substantial differentiation at an evolutionary timescale (Formia et al., 2006).

Our study further expands the geographic coverage of previous MSAs of the green turtle in the Atlantic, incorporating 14 nesting populations and 17 foraging grounds in our dataset. In particular the inclusion of African foraging grounds (i.e. Corisco Bay, Sao Tome and 'West Africa') improved the estimates for the distribution of hatchlings from Poilão, significantly reducing the estimate of the putative "unknown" foraging site (here 3.7%) compared to a recent MSA (14.3% in Putman and Naro-Maciel, 2013), as well as substantially reducing the confidence intervals. In a previous m2m MSA a high contribution of Ascension Island to Corisco Bay was estimated (ca. 40%, Bolker et al., 2007). Here that contribution drops to 9.2%, and we predict a much stronger connectivity between Poilão and Corisco. By including more foraging grounds in our analyses, we show that Ascension rookery contributes primarily to juvenile aggregations along the Southwest Atlantic (71.6%), also seen in Putman and Naro-Maciel (2013). Analogously, the foraging ground-centric MSA in Bolker et al (2007) attributes most of the Corisco Bay foraging ground to Ascension Island (>70%), while we estimate that 60.5% of the aggregation origins at Poilão, and only 27.7% would come from Ascension. Additionally, the contributions of Aves Island and NE Brazil to Corisco Bay estimated before (ca. 15% each, Bolker et al., 2007) were considerably lower in our study (2.7% and 4.8%, respectively), and these populations also seem to contribute more to the Southwest Atlantic. See tables S3 and S4 for m2m MSA summary results.

326

327 **Limitations of MSA and future directions**

328 Although increasing the available sample size at Poilão and expanding the dataset for
329 Atlantic green turtles has improved MSA estimates, this analysis is based on a single marker
330 and on a short fragment of the mtDNA. To further unveil the green turtle connectivity puzzle
331 in the Atlantic (and elsewhere) the strength of the genetic signal can be enhanced, at a
332 lesser cost than substantially increasing sample sizes. Data from the longer mtDNA
333 sequences should be obtained from existing samples and made available, to be
334 incorporated in MSAs. Additionally, a new marker consisting of four AT short tandem repeats
335 (STRs) in the 3' end of the mtDNA, the mtSTR, had been shown to add information on the
336 genetic variability within unique mtDNA haplotype classes and to contribute to improve the
337 knowledge on population connectivity and evolutionary relationships (Shamblin et al., 2015,
338 Tikochinski et al., 2012). Recent research using nuclear markers have found significant
339 structure among sea turtle rookeries, supportive of male philopatry (Carreras et al., 2011;
340 Naro-Maciel et al., 2012; Naro-Maciel et al., 2014; Roden et al., 2013). Finally, new genomic
341 approaches are have the potential to greatly increase the signal resolution and detect fine-
342 scale population structure (Benestan et al., 2015; Funk et al., 2012; Milano et al., 2014).
343 Some of the above information is now becoming available at local scales. Hopefully future
344 collaborations among research groups at wider scales will lead to significant advances in our
345 understanding of the dispersal and distribution of marine turtles.

346

347 **Adult linkage**

348 Godley et al (2010) recorded the trajectories of eight post-nesting females from Poilão using
349 satellite transmitters, finding that they foraged either locally, at the Bijagós Archipelago (n =
350 4) , or regionally (n = 4), at the Banc d'Arguin National Park, Mauritania (> 1000 km distant).
351 This aspect of investigation would clearly benefit from enhanced sampling effort, preferably
352 across multiple seasons, at different points of the season and across a range of size
353 classes, to avoid inter-annual (Witt et al., 2011), seasonal (Rees et al., 2010) and phenotypic

(Hawkes et al., 2006) biases in dispersal. Future satellite tracking should be conducted in tandem with stable isotope analysis to facilitate the posterior assignment of turtles to these areas, facilitating the analyses of larger sample sizes, more relevant for population studies (Zbinden et al., 2011).

If nesting females from Poilão are limited to the East Atlantic it does not necessarily contradict our suggestion of transatlantic dispersal as post-hatchlings. Post-hatchling turtles forage during their developmental migration (Reich et al., 2007), which allows them to travel much longer distances than adults that typically fast during their reproductive migrations (Hays and Scott, 2013; Scott et al., 2014). According to Scott et al (2014), if the developmental foraging area is so far as to be too costly to be repeatable during the cyclic reproductive migrations, adults may forage locally, as observed at the Bijagós, instead of returning to the sites experienced when younger. This mechanism reduces the consumption of reproductive energy utilized, potentially increasing fecundity, however it is dependent on the availability of foraging areas.

Conservation implications

In this study we show the importance of Poilão rookery for the recruitment of juvenile green turtles in West Africa, and also that the link with the Southwest Atlantic is very likely. In Guinea-Bissau, despite marine turtles being fully protected by the national fisheries law, illegal take continues to occur without much law enforcement effort (Catry et al., 2009), particularly at the Bijagós Archipelago, where turtles are frequently harvested at the nesting beaches, mostly for local consumption (Catry et al., 2009). The nesting population at Poilão is one exception, thanks to the Bijagós traditional 'law' (reinforced by state authorities), restricting access to the island on very rare ceremonies of social and religious significance (Catry et al., 2009). Off Guinea-Bissau and along the coast of West Africa however, vast artisanal fleets and many industrial fishing fleets operate, using trawlers without turtle excluder devices (Catry et al., 2009; Zeeberg et al., 2006), and longlining (Moore et al.,

2010). Unfortunately there is a scarcity of quantitative data in the region, either on bycatch or on targeted harvesting of marine turtles, particularly from artisanal fisheries (Moore et al., 2010). The foraging grounds in the Southwest Atlantic to which Poilão seems to contribute to, on the other hand, are mostly protected from illegal harvesting (Marcovaldi and dei Marcovaldi, 1999), although bycatch may be a problem (Wallace et al., 2010). Despite the existing threats, major green turtle populations are recovering globally following decades of conservation efforts (Bourjea et al., 2015, Broderick et al., 2006, Catry et al., 2009). It may be that the long term enhanced protection in South America and the efforts in Poilão itself are the principle factors involved in the recovery of this population.

Acknowledgments

We thank the Institute of Biodiversity and Protected Areas of Guinea-Bissau (IBAP-GB) for all the logistic support for sample collection and all the people involved in the field-work, particularly the community members from the Bijagós, and the rangers and technicians from the IBAP. Sampling permits were obtained by the IBAP-GB, CITES export license was obtained from the Directorate General of Forest and Fauna of Guinea-Bissau (DGFF-GB), and CITES import license (13-PT-LX0006/P) was emitted by the Institute for Nature Conservation and Forests (ICNF-PT). Research was conducted with the financial support from the MAVA Foundation, the Rufford Foundation (RSG12317-1, RSG16357-2), and the Portuguese Foundation for Science and Technology through the strategic project UID/MAR/04292/2013 granted to MARE, project IF/00502/2013/CP1186/CT0003 and the grant awarded to ARP (fellowship SFRH/BD/85017/2012). BJG was supported by the Darwin Initiative.

405 **Literature cited**

- 406 Abreu-Grobois A, Horrock J, Formia A, Dutton P, LeRoux R, Velez-Zuazo X, Soares L and
 407 Meylan P (2006) New mtDNA Dloop primers which work for a variety of marine turtles
 408 species may increase the resolution of mixed stock analyses. In: Frick M,
 409 Panagopoulou A, Rees A, Williams K (eds). Proc 26th Annual Symp Sea Turtle Biology
 410 and Conservation, International Sea Turtle Society. Island of Crete, Greece, p 179
 411 Bagley D (2003) Characterizing juvenile green turtles, (*Chelonia mydas*), from three east
 412 central Florida developmental habitats. Master's dissertation, University of Central
 413 Florida, Orlando, FL
 414 Bass AL, Epperly SP, Braun-McNeill J (2006) Green Turtle (*Chelonia mydas*) Foraging and
 415 Nesting Aggregations in the Caribbean and Atlantic: Impact of Currents and Behavior
 416 on Dispersal. J Hered 97: 346–354 doi:10.1093/jhered/esl004
 417 Bass AL, Witzell WN (2000) Demographic composition of immature green turtles (*Chelonia*
 418 *mydas*) from the east central Florida coast: evidence from mtDNA markers.
 419 Herpetologica 56: 357–367
 420 Bjørndal K, Bolten A, Trøeng S (2005) Population structure and genetic diversity in green
 421 turtles nesting at Tortuguero, Costa Rica, based on mitochondrial DNA control region
 422 sequences. Mar Biol 147: 1449–1457. doi:10.1007/s00227-005-0045-y
 423 Bjørndal KA, Bolten AB, Moreira L, Bellini C, Marcovaldi MÂ (2006) Population Structure and
 424 Diversity of Brazilian Green Turtle Rookeries Based on Mitochondrial DNA Sequences.
 425 Chelonian Conserv Biol 5: 262–268 doi:10.2744/1071-
 426 8443(2006)5[262:psadob]2.0.co;2
 427 Bolker B, Okuyama T, Bjørndal KA, Bolten AB (2007) Incorporating multiple mixed stocks in
 428 mixed stock analysis: “Many-to-many” analyses. Mol Ecol 16: 685–695.
 429 doi:10.1111/j.1365-294X.2006.03161.x
 430 Bolten AB (2003) Variation in sea turtle life history patterns: neritic versus oceanic
 431 developmental stages, in: Lutz, P.L., Musick, J.A., Wyneken, J. (eds.), The Biology of
 432 Sea Turtles. CRC Press, Boca Raton, Florida, pp. 243–257
 433 Bonfil R, Meyer M, Scholl MC, Johnson R, O'Brien S, Oosthuizen H, Swanson S, Kotze D,
 434 Paterson M (2005) Transoceanic migration, spatial dynamics, and population linkages
 435 of white sharks. Science 310: 100–103 doi:10.1126/science.1114898
 436 Booth DT (2014) Kinematics of swimming and thrust production during powerstroking bouts
 437 of the swim frenzy in green turtle hatchlings. Biol Open 3: 887–894.
 438 doi:10.1242/bio.20149480
 439 Bourjea J, Dalleau M, Derville S, Beudard F, Marmoeux C, M'Soili A, Roos D, Ciccione S and
 440 Frazier J (2015) Seasonality, abundance, and fifteen-year trend in green turtle nesting
 441 activity at Itsamia, Moheli, Comoros. Endang Species Res 27: 265-276
 442 Bowen BW, Karl SA (2007) Population genetics and phylogeography of sea turtles. Mol Ecol
 443 16: 4886–4907 doi:10.1111/j.1365-294X.2007.03542.x
 444 Broderick AC, Frauenstein R, Glen F, Hays GC, Jackson AL, Pelembe T, Ruxton GD and
 445 Godley BJ (2006). Are green turtles globally endangered? Global Ecol Biogeogr 15: 21-
 446 26
 447 Carreras C, Pascual M, Cardona L, Marco A, Bellido JJ, Castillo JJ, Tomás J, Raga JA,
 448 Sanfélix M, Fernández G and Aguilar A (2011) Living together but remaining apart:
 449 Atlantic and Mediterranean loggerhead sea turtles (*Caretta caretta*) in shared feeding
 450 grounds. J Hered 102: 666–677 doi:10.1093/jhered/esr089
 451 Catry P, Barbosa C, Indjai B, Almeida A, Godley BJ, Vié JC (2002) First census of the green
 452 turtle at Poilão, Bijagós Archipelago, Guinea-Bissau: the most important nesting colony
 453 on the Atlantic coast of Africa. Oryx 36: 400–403 doi:doi:10.1017/S0030605302000765
 454 Catry P, Barbosa C, Paris B, Indjai B, Almeida A (2009) Status, Ecology, and Conservation
 455 of Sea Turtles in Guinea-Bissau 8:150–160
 456 Catry P, Dias MP, Phillips RA, Granadeiro JP (2011) Different means to the same end:
 457 Long-distance migrant seabirds from two colonies differ in behaviour, despite common
 458 wintering grounds. PLoS One 6: 4–9 doi:10.1371/journal.pone.0026079

- Encalada SE, Lahanas PN, Bjorndal KA, Bolten AB, Miyamoto MM, Bowen BW, (1996) Phylogeography and population structure of the Atlantic and Mediterranean green turtle *Chelonia mydas*: a mitochondrial DNA control region sequence assessment. *Mol Ecol* 5: 473–483 doi:10.1111/j.1365-294X.1996.tb00340.x
- Excoffier L, Lischer HE (2010) Arlequin suite ver 3.5: a new series of programs to perform population genetics analyses under Linux and Windows. *Molecular ecology resources*, 10: 564–567
- Formia A (2002) Population and genetic structure of the green turtle (*Chelonia mydas*) in West and Central Africa: implications for management and conservation. PhD dissertation, Cardiff University, Cardiff, UK
- Formia A, Godley BJ, Dontaine JF, Bruford MW (2006) Mitochondrial DNA diversity and phylogeography of endangered green turtle (*Chelonia mydas*) populations in Africa. *Conserv Genet* 7: 353–369 doi:10.1007/s10592-005-9047-z
- Formia A, Broderick AC, Glen F, Godley BJ, Hays GC, Bruford MW (2007). Genetic composition of the Ascension Island green turtle rookery based on mitochondrial DNA: implications for sampling and diversity. *Endang Species Res* 3: 145–158
- Fukuoka T, Narazaki T and Sato K (2015) Summer-restricted migration of green turtles *Chelonia mydas* to a temperate habitat of the northwest Pacific Ocean. *Endang Species Res* 28: 1 – 10
- Godley BJ, Barbosa C, Bruford M, Broderick AC, Catry P, Coyne MS, Formia A, Hays GC, Witt MJ (2010) Unravelling migratory connectivity in marine turtles using multiple methods. *J Appl Ecol* 47: 769–778 doi:10.1111/j.1365-2664.2010.01817.x
- Hall TA (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. In *Nucleic acids symposium series* 41:95-98
- Hawkes LA, Broderick AC, Coyne MS, Godfrey MH, Lopez-Jurado LF, Lopez-Suarez P, Merino SE, Varo-Cruz N, Godley BJ (2006). Phenotypically linked dichotomy in sea turtle foraging requires multiple conservation approaches. *Curr Biol* 16: 990–995
- Hays GC, Scott R (2013) Global patterns for upper ceilings on migration distance in sea turtles and comparisons with fish, birds and mammals. *Funct Ecol* 27: 748–756 doi:10.1111/1365-2435.12073
- Lahanas PN, Bjorndal KA, Bolten AB, Encalada SE, Miyamoto MM, Valverde RA, Bowen BW (1998) Genetic composition of a green turtle (*Chelonia mydas*) feeding ground population: evidence for multiple origins. *Mar Biol* 130: 345–352 doi:10.1007/s002270050254
- Lahanas PN, Miyamoto MM, Bjorndal KA, Bolten AB (1994) Molecular evolution and population genetics of Greater Caribbean green turtles (*Chelonia mydas*) as inferred from mitochondrial DNA control region sequences. *Genetica* 94: 57–66 doi:10.1007/BF01429220
- Luke K, Horrocks JA, LeRoux RA, Dutton PH (2004). Origins of green turtle (*Chelonia mydas*) feeding aggregations around Barbados, West Indies. *Mar Biol* 144: 799–805 doi:10.1007/s00227-003-1241-2
- Lunn D, Thomas A, Best N, Spiegelhalter D (2000) WinBUGS – A Bayesian modelling framework: Concepts, structure, and extensibility. *Stat Comput* 10: 325–337 doi:10.1023/A:1008929526011
- Marcovaldi MÂ, Dei Marcovaldi GG (1999) Marine turtles of Brazil: the history and structure of Projeto TAMAR-IBAMA. *Biol Conserv* 91: 35–41 doi:10.1016/S0006-3207(99)00043-9
- Meylan AB, Bowen BW, Avise JC (1990) A genetic test of the natal homing versus social facilitation models for green turtle migration. *Science* 248: 724–727 doi:10.1126/science.2333522
- Millar RB (1987) Maximum likelihood estimation of mixed stock fishery composition. *Can J Fish Aquat Sci* 44: 583–590
- Monzón-Argüello C, López-Jurado LF, Rico C, Marco A, López P, Hays GC, Lee PLM (2010) Evidence from genetic and Lagrangian drifter data for transatlantic transport of small juvenile green turtles. *J Biogeogr* 37: 1752–1766 doi:10.1111/j.1365-

2699.2010.02326.x

Moore JE, Cox TM, Lewison RL, Read AJ, Bjorkland R, McDonald SL, Crowder LB, Aruna E, Ayissi I, Espeut P, Joynson-Hicks C, Pilcher N, Poonian CNS, Solarin B, Kiszka J (2010) An interview-based approach to assess marine mammal and sea turtle captures in artisanal fisheries. *Biol Conserv* 143: 795–805 doi:10.1016/j.biocon.2009.12.023

Naro-Maciel E, Becker JH, Lima EHSM, Marcovaldi MÂ, DeSalle R (2007) Testing Dispersal Hypotheses in Foraging Green Sea Turtles (*Chelonia mydas*) of Brazil. *J Hered* 98: 29–39 doi:10.1093/jhered/esl050

Naro-Maciel E, Bondioli A, Martin M, De Pádua Almeida A, Baptistotte C, Bellini C, Marcovaldi MÂ, Santos AJB, Amato G (2012) The interplay of homing and dispersal in green turtles: A focus on the southwestern atlantic. *J Hered* 103: 792–805 doi:10.1093/jhered/ess068

Naro-Maciel E, Reid BN, Alter SE, Amato G, Bjorndal KA, Bolten AB, Martin M, Nairn CJ, Shamblin B, Pineda-catalan O (2014) From refugia to rookeries : Phylogeography of Atlantic green turtles. *J Exp Mar Bio Ecol* 461: 306–316

Naro-Maciel E, Hart KM, Cruciata R, Putman NF (2016) DNA and dispersal models highlight constrained connectivity in a migratory marine megavertebrate. *Ecography* doi:10.1111/oik.03181

Narum SR (2006) Beyond Bonferroni: Less conservative analyses for conservation genetics. *Conserv Genet* 7: 783–787 doi:10.1007/s10592-005-9056-y

Patrício AR, Velez-Zuazo X, Diez CE, van Dam R, Sabat AM (2011) Survival probability of immature green turtles in two foraging grounds at Culebra, Puerto Rico. *Mar Ecol Prog Ser* 440: 217–227

Patrício R, Diez CE and van Dam RP (2014) Spatial and temporal variability of immature green turtle abundance and somatic growth in Puerto Rico. *Endang Species Res* 23: 51-62

Peakall R, Smouse PE (2012) GenALEX 6.5: Genetic analysis in Excel. Population genetic software for teaching and research-an update. *Bioinformatics* 28: 2537–2539 doi:10.1093/bioinformatics/bts460

Pella J, Masuda M (2001) Bayesian methods for analysis of stock mixtures from genetic characters. *Fish Bull* 99: 151–167

Proietti M, Reisser JW, Kinas PG, Kerr R, Monteiro DS, Marins LF, Secchi ER (2012) Green turtle *Chelonia mydas* mixed stocks in the western South Atlantic, as revealed by mtDNA haplotypes and drifter trajectories. *Mar Ecol Prog Ser* 447: 195–209 doi:10.3354/meps09477

Prosdocimi L, González Carman V, Albareda DA, Remis MI (2012) Genetic composition of green turtle feeding grounds in coastal waters of Argentina based on mitochondrial DNA. *J Exp Mar Bio Ecol* 412: 37–45 doi:10.1016/j.jembe.2011.10.015

Putman N, Scott R, Verley P, Marsh R, Hays GC (2012a) Natal site and offshore swimming influence fitness and long-distance ocean transport in young sea turtles. *Mar Biol* 159: 2117–2126 doi:10.1007/s00227-012-1995-5

Putman N, Verley P, Shay TJ, Lohmann KJ (2012b) Simulating transoceanic migrations of young loggerhead sea turtles: merging magnetic navigation behavior with an ocean circulation model. *J Exp Biol* 215: 1863–1870 doi:10.1242/jeb.067587

Putman N, Naro-maciel E (2013) Finding the “lost years” in green turtles: insights from ocean circulation models and genetic analysis. *Proc R Soc London Ser B* 280: 20131468 doi:10.1098/rspb.2013.1468

Putman NF, Mansfield KL (2015) Direct Evidence of Swimming Demonstrates Active Dispersal in the Sea Turtle “Lost Years.” *Curr Biol* 25: 1221-1227 doi:10.1016/j.cub.2015.03.014

Rasmussen K, Palacios DM, Calambokidis J, Saborío MT, Dalla Rosa L, Secchi ER, Steiger GH, Allen JM, Stone GS (2007) Southern Hemisphere humpback whales wintering off Central America: insights from water temperature into the longest mammalian migration. *Biol Lett* 3: 302–305 doi:10.1098/rsbl.2007.0067

Rees AF, Saady SA, Broderick AC, Coyne MS, Papathanasopoulou N, Godley BJ (2010)


- Behavioural polymorphism in one of the world's largest populations of loggerhead sea turtles *Caretta caretta*. *Mar Ecol Prog Ser* 418: 201-12
- Rees AF, Alfaro-Shigueto J, Barata PCR, Bjorndal KA, Bolten AB, Bourjea J, Broderick AC, Campbell LM, Cardona L, Carreras C, Casale P, Ceriani SA, Dutton PH, Eguchi T, Formia A, Fuentes MMPB, Fuller WJ, Girondot M, Godfrey MH, Hamann M, Hart KM, Hays GC, Hochscheid S, Kaska Y, Jensen MP, Mangel JC, Mortimer JA, Naro-Maciel E, Ng CKY, Nichols WJ, Phillott AD, Reina RD, Revuelta O, Schofield G, Seminoff JA, Shanker K, Tomás J, van de Merwe JP, Van Houtan KS, Vander Zanden HB, Wallace BP, Wedemeyer-Strombel KR, Work TM, Godley BJ (2016) Are we working towards global research priorities for management and conservation of sea turtles? *Endang Spec Res* 31:337-382
- Reich KJ, Bjorndal KA, Bolten, A.B., 2007. The "lost years" of green turtles: using stable isotopes to study cryptic lifestages. *Biol Lett* 3: 712–714 doi:10.1098/rsbl.2007.0394
- Roden SE, Morin PA, Frey A, Balazs GH, Zarate P, Cheng IJ, Dutton PH (2013) Green turtle population structure in the Pacific: New insights from single nucleotide polymorphisms and microsatellites. *Endang Species Res* 20: 227–234 doi:10.3354/esr00500
- Rooker JR, Arrizabalaga H, Fraile I, Secor DH, Dettman DL, Abid N, Addis P, Deguara S, Karakulak FS, Kimoto A, Sakai O, Macías D, Santos MN (2014) Crossing the line: Migratory and homing behaviors of Atlantic bluefin tuna. *Mar Ecol Prog Ser* 504: 265–276 doi:10.3354/meps10781
- Ruiz-Urquiola A, Riverón-Giró FB, Pérez-Bermúdez E, Abreu-Grobois FA, González-Pumariiega M, James-Petric BL, Díaz-Fernández R, Álvarez-Castro JM, Jager M, Azanza Ricardo J, Espinosa-López G (2010) Population genetic structure of greater Caribbean green turtles (*Chelonia mydas*) based on mitochondrial DNA sequences, with an emphasis on rookeries from southwestern Cuba. *Rev Investig Mar* 31: 33–52
- Scott R, Marsh R, Hays GC (2012) A little movement orientated to the geomagnetic field makes a big difference in strong flows. *Mar Biol* 159: 481–488 doi:10.1007/s00227-011-1825-1
- Scott R, Marsh R, Hays G (2014) Ontogeny of long distance migration. *Ecology* 95: 2840–2850 doi:10.1890/13-2164.1
- Seminoff JA, Allen CD, Balazs GH, Dutton PH, Eguchi T, Haas HL, Hargrove SA, Jensen MP, Klemm DL, Lauritsen AM, MacPherson SL, Opat P, Possardt EE, Pultz SL, Seney EE, Van Houtan KS, Waples RS (2015) Status Review of the Green Turtle (*Chelonia mydas*) Under the U.S. Endangered Species Act. NOAA Technical Memorandum, NOAA NMFS- SWFSC-539
- Shamblin BM, Bjorndal KA, Bolten AB, Hillis-Starr ZM, Lundgren IAN, Naro-Maciel E, Nairn CJ (2012) Mitogenomic sequences better resolve stock structure of southern Greater Caribbean green turtle rookeries. *Mol Ecol* 21: 2330–2340 doi:10.1111/j.1365-294X.2012.05530.x
- Shamblin BM, Bagley DA, Ehrhart LM, Desjardin NA, Martin RE, Hart KM, Naro-Maciel E., Rusenko K, Stiner JC, Sobel D, Johnson C, Wilmers TJ, Wright LJ, Nairn CJ (2014) Genetic structure of Florida green turtle rookeries as indicated by mitochondrial DNA control region sequences. *Conserv Genet* 16: 673–685 doi:10.1007/s10592-014-0692-y
- Shamblin BM, Dutton PH, Bjorndal KA, Bolten AB, Naro-Maciel E, Santos AJB, Bellini C, Baptistotte C, Marcovaldi MÂ and Nairn CJ (2015) Deeper Mitochondrial Sequencing Reveals Cryptic Diversity and Structure in Brazilian Green Turtle Rookeries. *Chelonian Conserv Biol* 14: 167-172
- Shimada T, Aoki S, Kameda K, Hazel J, Reich K and Kamezaki N (2014) Site fidelity, ontogenetic shift and diet composition of green turtles *Chelonia mydas* in Japan inferred from stable isotope analysis. *Endang Species Res* 25: 151-164
- Tikochinski Y, Bendelac R, Barash A, Daya A, Levy Y, Friedmann A (2012) Mitochondrial DNA STR analysis as a tool for studying the green sea turtle (*Chelonia mydas*) populations: The Mediterranean Sea case study. *Mar Genomics* 6: 17–24 doi:10.1016/j.margen.2012.01.002
- Velez-Zuazo X, Ramos WD, van Dam RP, Diez CE, Abreu-Grobois A, McMillan WO (2008)

624 Dispersal, recruitment and migratory behaviour in a hawksbill sea turtle aggregation.
 625 Mol Ecol 17: 839–853 doi:10.1111/j.1365-294X.2007.03635.x
 626 Wallace BP, Lewison RL, McDonald SL, McDonald RK, Kot CY, Kelez S, Bjorkland RK,
 627 Finkbeiner EM, Helmbrecht S, Crowder LB (2010) Global patterns of marine turtle
 628 bycatch. Conserv Lett 3: 131–142 doi:10.1111/j.1755-263X.2010.00105.x
 629 Witt MJ, Bonguno EA, Broderick AC, Coyne MS, Formia A, Gibudi A, Mounquengui GA,
 630 Moussounda C, NSafou M, Nougessono S, Parnell RJ (2011) Tracking leatherback
 631 turtles from the world's largest rookery: assessing threats across the South Atlantic.
 632 Proc R Soc Lond B Biol Sci 278: 2338-2347
 633 Zbinden JA, Bearhop S, Bradshaw P, Gill B, Margaritoulis D, Newton J, Godley BJ, (2011)
 634 Migratory dichotomy and associated phenotypic variation in marine turtles revealed by
 635 satellite tracking and stable isotope analysis. Mar Ecol Prog Ser 421: 291–302
 636 doi:10.3354/meps08871
 637 Zeeberg J, Corten A, de Graaf E (2006) Bycatch and release of pelagic megafauna in
 638 industrial trawler fisheries off Northwest Africa. Fish Res 78: 186–195
 639 doi:10.1016/j.fishres.2006.01.012

640 **Table 1.** Haplotypic and genetic diversity of Atlantic green turtle (*Chelonia mydas*) nesting populations (n = 14) included in a many-to-many
641 mixed stock analysis (Bolker et al., 2007), using the control region of the mitochondrial DNA as a marker (490bp). The number of females
642 refers to the total number of reproductive females in each population. The study population is represented in bold.

Nesting Population	short name	sample size	no. of females*	no. of haplotypes	haplotypic diversity (<i>h</i>)	nucleotide diversity (π)
East central Florida	EcFL	311	4490	9	0.512 \pm 0.020	0.0016 \pm 0.0013
South Florida	SFL	174	3302	10	0.444 \pm 0.043	0.0022 \pm 0.0016
Southwest Cuba	CUB	28	2226	7	0.648 \pm 0.089	0.0053 \pm 0.0033
Quintana Roo, Mexico	MEX	20	18257	7	0.816 \pm 0.058	0.0051 \pm 0.0032
Tortuguero, Costa Rica	CR	433	131751	5	0.163 \pm 0.023	0.0033 \pm 0.0022
Matapica/Galibi, Suriname	SUR	73	13067	4	0.132 \pm 0.053	0.0013 \pm 0.0011
Buck Island	BUC	49	63	2	0.153 \pm 0.065	0.0030 \pm 0.0020
Aves Island	AV	67	2833	2	0.140 \pm 0.055	0.0029 \pm 0.0020
Rocas/Fernando Noronha	RC/FN	69	345	7	0.463 \pm 0.071	0.0026 \pm 0.0018
Trindade Island	TRI	99	2016	7	0.505 \pm 0.052	0.0012 \pm 0.0011
Ascension Island	ASC	245	1417	13	0.303 \pm 0.038	0.0008 \pm 0.0008
Poilão, Guinea-Bissau	GB	171	29016	2	0.012 \pm 0.011	0.0001 \pm 0.0003
Bioko Island, Eq. Guinea	BIO	50	850	2	0.184 \pm 0.068	0.0004 \pm 0.0006
Sao Tome and Principe	STP	26	376	7	0.569 \pm 0.110	0.0026 \pm 0.0019

643
644 * Seminoff et al., (2015)



650 2005; Encalada et al., 1996); SUR: Matapica and Galibi, Suriname (Encalada et al., 1996; Shamblin et al., 2012); AV: Aves Island (Lahanas et
 651 al., 1998, 1994; Shamblin et al., 2012), Venezuela; BUC: Buck Island (Shamblin et al., 2012); RC/FN: Rocas Atol and Fernando de Noronha
 652 (Bjorndal et al., 2006; Encalada et al., 1996), Brazil; ASC: Ascension Island (Encalada et al., 1996; Formia et al., 2007); TRI: Trindade Island,
 653 Brazil (Bjorndal et al., 2006); POI: Poilão, Guinea-Bissau (study population); BIO: Bioko Island, Equatorial Guinea (Formia et al., 2006); STP:
 654 Sao Tome and Principe (Formia et al., 2006). **Foraging grounds:** NC: North Carolina (Bass et al., 2006), EcFL: East central Florida (Bagley,
 655 2003; Bass and Witzell, 2000), BHM: Bahamas (Lahanas et al., 1998), BRB: Barbados (Luke et al., 2004), ALF: Almofala, Brazil (Naro-Maciel
 656 et al., 2007), RC: Rocas Atoll, Brazil (Naro-Maciel et al., 2012), FN: Fernando de Noronha, Brazil (Naro-Maciel et al., 2012), BA: Bahia, Brazil
 657 (Naro-Maciel et al., 2012), ES: Espirito Santo, Brazil (Naro-Maciel et al., 2012), UB: Ubatuba, Brazil (Naro-Maciel et al., 2007), AI: Arvoredo
 658 Island, Brazil (Proietti et al., 2012), CB: Cassino Beach, Brazil (Proietti et al., 2012), BuA, Buenos Aires, Argentina (Prosdocimi et al., 2012),
 659 CV: Cape Verde (Monzón-Argüello et al., 2010), COR: Corisco Bay, Equatorial Guinea (Formia, 2002), WA: 'West Africa' (Liberia to Benin,
 660 Formia, 2006). *GfC*: Gulf Current, *NEC*: North Equatorial Current, *SEC*: South Equatorial Current, *BrC*: Brazil Current, *GC*: Guinea Current,
 661 *BgC*: Benguela Current; **b) Region map** with study site, Poilão, and three juvenile foraging grounds likely to partly originate at Poilão, to date
 662 genetically uncharacterized: Unhocomo/Unhocomozinho and Varela at Guinea Bissau, and Banc d'Arguin at Mauritania. Dashed arrow
 663 illustrates the direction of four green turtle females tracked from Poilão to Banc d'Arguin (Godley et al., 2010). (Maps created using
 664 www.seaturtle.org/maptool).





Figure 2. PCoA of the 14 Atlantic green turtle (*Chelonia mydas*) populations using ϕ_{st} and considering the 490bp fragment. The percentage of the variability explained by each coordinate is shown in brackets. ASC: Ascension Island; AV: Aves Island, Venezuela; BIO: Bioko Island, Equatorial Guinea; BUC: Buck Island; CR: Tortuguero, Costa Rica; CUB: Southwest Cuba; EcFL: East central Florida, MEX: Quintana Roo, Mexico; POI: Poilão, Guinea-Bissau (this study); RC/FN: Rocas Atol and Fernando Noronha, Brazil; SFL: South Florida, USA; STP: Sao Tome and Principe; SUR: Matapica and Galibi, Suriname; TRI: Trindade Island, Brazil.



675

676

677

678

679

680

681

682

Figure 3. Mean relative contribution of Poilão (Guinea-Bissau) nesting population to 17 Atlantic green turtle (*Chelonia mydas*) foraging grounds, estimated by a many2many mixed stock analysis (Bolker et al 2007). Error bars show 95 % confidence intervals. WA: 'West Africa' – Liberia to Benin, ST: Sao Tome, COR: Corisco Bay, CV: Cape Verde, BuA: Buenos Aires, UB: Ubatuba, ALF: Almofala, CB: Cassino Beach, FN: Fernando de Noronha, ES: Espírito Santo, BA: Bahia, AI: Arvoredo Island, RC: Rocas Atol, BRB: Barbados, BHM: Bahamas, NC: North Carolina, EcFL: East central Florida. Dashed lines separate geographic regions.

Supplementary material:

Table S1. mtDNA control region haplotype frequencies (490 bp), at 14 Atlantic green turtle nesting populations with total no. of samples per area. See Fig. 1 for site abbreviations. Long haplotypes (856bp) for study area are shown in the table below.

Haplotype	Nesting Populations												
	EcFL ^{a,b}	SFL ^b	MEX ^a	CR ^{c,d}	CUB ^e	BUC ^f	AV ^{d,f,g}	SUR ^{a,f}	RC/N ^{a,h}	TRI ^h	ASC ^{a,i,j}	POI ^k	BIO ⁱ
CM-A1	197	27	7		3								
CM-A2	7	4											
CM-A3	92	127	5	395	16		5	1					
CM-A4				1		16							
CM-A5	2	4	1	32		45	62	68					
CM-A6								3			11		5
CM-A7								1					
CM-A8	1								50	67	204	170*	45
CM-A9									7	19	9		
CM-A10									2		5		
CM-A11									1	1			
CM-A12									5				
CM-A13	7	2											
CM-A15			1										
CM-A16	2	1	1										
CM-A17		2	2										
CM-A18	1	1	3										
CM-A20				2									
CM-A21				3									
CM-A23										6	1		
CM-A24										1	7		
CM-A25									3		1		
CM-A27					1								
CM-A28	2	3			1								
CM-A32									1	4	1		
CM-A33										1			
CM-A35													
CM-A36													
CM-A37													
CM-A38													
CM-A39											1		
CM-A42												1*	
CM-A44											1		
CM-A45											1		
CM-A46											2		
CM-A48					5								
CM-A50											1		
CM-A53		3											
CM-A56					1								
CM-A57					1								
n	311	174	20	433	28	61	67	73	69	99	245	171	50

^aEncalada et al. 1996, ^bShamblin et al. 2014, ^cBjorndal et al. 2005, ^dLahanas et al. 1998, ^eRuiz-Urquiola et al. 2010, ^fShamblin et al. 2012, ^gLahanas et al. 1994, ^hBjorndal et al. 2006, ⁱFormia et al. 2006, ^jFormia et al. 2007, ^kThis study

* Long haplotypes (856bp): CMA8.1 (n=169), CMA8.3 (n=1), CMA42.1 (n=1)

Table S2. Pairwise exact test P-values (above diagonal) and pairwise F_{ST} values (below diagonal) among the 14 Atlantic green turtle (*Chelonia mydas*) nesting populations, based on ~490bp sequences of the control region of the mtDNA. The study site is in grey and in bold, and abbreviations follow those in Figure 1. Asterisks indicate statistically significant comparisons (* $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$) i) prior to corrections, in the low diagonal, ii) after false discovery rate (FDR) correction, in the above diagonal. Non-significant values, after FDR (Narum, 2006) correction, are marked in bold (for a $P < 0.05$ FDR = 0.0098, $P < 0.01$ FDR = 0.0020, $P < 0.001$ FDR = 0.0002).

	MEX	EcFL	SFL	CR	AV	BUC	CUB	SUR	TRI	RC/FN	ASC	POI	BIO	STP
MEX	-	0.009*	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***
EcFL	0.082**	-	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***
SFL	0.182***	0.197***	-	0.000***	0.000***	0.000***	0.009*	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***
CR	0.202***	0.254***	0.033***	-	0.000***	0.000***	0.009*	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***
AV	0.796***	0.895***	0.872***	0.820***	-	0.342	0.000***	0.108	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***
BUC	0.783***	0.897***	0.873***	0.822***	0.000	-	0.000***	0.045	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***
CUB	0.104***	0.243***	0.131**	0.154***	0.822***	0.811***	-	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***
SUR	0.880***	0.920***	0.905***	0.849***	0.021	0.031*	0.887***	-	0.000***	0.000***	0.000***	0.000***	0.000***	0.000***
TRI	0.860***	0.899***	0.885***	0.820***	0.657***	0.659***	0.873***	0.759***	-	0.009*	0.000***	0.000***	0.000***	0.000***
RC/FN	0.787***	0.886***	0.863***	0.810***	0.567***	0.554***	0.812***	0.666***	0.031**	-	0.000***	0.000***	0.000***	0.009*
ASC	0.913***	0.918***	0.914***	0.852***	0.728***	0.735***	0.922***	0.795***	0.060***	0.037***	-	0.000***	0.243	0.000***
POI	0.953***	0.931***	0.929***	0.855***	0.805***	0.823***	0.950***	0.895***	0.146***	0.070***	0.016***	-	0.000***	0.000***
BIO	0.877***	0.909***	0.894***	0.824***	0.640***	0.646***	0.878***	0.789***	0.093***	0.037***	0.003	0.106***	-	0.036
STP	0.766***	0.895***	0.870***	0.811***	0.522***	0.505***	0.792***	0.671***	0.083***	0.036*	0.067***	0.201***	0.045*	-

Table S3. Summary of source-centric mixed stock analysis of Atlantic green turtle nesting populations (n=14) and juvenile foraging grounds (n=17).

Nesting Population	Foraging grounds																	
	NC	EcFL	BHM	BRB	ALF	RC	FN	BA	ES	UB	AI	CB	BuA	CV	COR	ST	WA	X
Poilão, Guinea Bissau																		
Mean	0.01	0.01	0.03	0.03	0.04	0.02	0.04	0.04	0.04	0.05	0.03	0.04	0.06	0.06	0.10	0.17	0.18	0.04
CI: 97.5%	0.04	0.05	0.09	0.10	0.10	0.06	0.09	0.10	0.09	0.14	0.08	0.10	0.13	0.13	0.20	0.31	0.32	0.13
CI: 2.5%	0.00	0.00	0.00	0.00	0.01	0.00	0.01	0.00	0.01	0.01	0.01	0.01	0.02	0.02	0.03	0.07	0.08	0.00
Bioko, Eq.Guinea																		
Mean	0.05	0.05	0.05	0.05	0.05	0.06	0.05	0.06	0.07	0.05	0.05	0.05	0.05	0.06	0.07	0.05	0.06	0.05
CI: 97.5%	0.17	0.17	0.19	0.19	0.18	0.20	0.19	0.20	0.22	0.19	0.19	0.18	0.19	0.20	0.22	0.19	0.21	0.18
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Sao Tome and Principe																		
Mean	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.05	0.06	0.05	0.06	0.09	0.08	0.06	0.05
CI: 97.5%	0.17	0.18	0.18	0.19	0.17	0.17	0.17	0.17	0.17	0.18	0.17	0.22	0.16	0.19	0.31	0.27	0.19	0.20
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Ascension Island, UK																		
Mean	0.01	0.01	0.02	0.05	0.09	0.09	0.05	0.05	0.08	0.11	0.09	0.08	0.07	0.02	0.09	0.03	0.04	0.02
CI: 97.5%	0.04	0.05	0.06	0.12	0.17	0.18	0.12	0.14	0.16	0.21	0.18	0.16	0.16	0.06	0.17	0.09	0.10	0.07
CI: 2.5%	0.00	0.00	0.00	0.00	0.03	0.03	0.01	0.00	0.02	0.04	0.03	0.02	0.02	0.00	0.03	0.00	0.00	0.00
Trindade, Brazil																		
Mean	0.03	0.03	0.04	0.04	0.04	0.08	0.04	0.12	0.08	0.05	0.10	0.09	0.08	0.03	0.03	0.04	0.03	0.04
CI: 97.5%	0.11	0.12	0.14	0.14	0.13	0.21	0.13	0.29	0.21	0.16	0.24	0.22	0.23	0.12	0.10	0.14	0.12	0.14
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00
Rocas/F.Noronha, Brazil																		
Mean	0.05	0.05	0.06	0.06	0.05	0.06	0.06	0.06	0.06	0.06	0.05	0.07	0.05	0.05	0.05	0.06	0.05	0.05
CI: 97.5%	0.20	0.19	0.19	0.20	0.19	0.22	0.19	0.19	0.20	0.20	0.19	0.23	0.18	0.19	0.16	0.19	0.18	0.19
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Suriname																		
Mean	0.01	0.03	0.05	0.07	0.07	0.05	0.14	0.07	0.08	0.03	0.04	0.04	0.05	0.17	0.01	0.02	0.03	0.04
low C.I.	0.04	0.09	0.15	0.16	0.15	0.11	0.25	0.15	0.15	0.07	0.09	0.09	0.12	0.31	0.03	0.06	0.09	0.12
upper C.I.	0.00	0.00	0.00	0.01	0.02	0.01	0.05	0.01	0.02	0.00	0.01	0.01	0.01	0.06	0.00	0.00	0.00	0.00

Table S3. Continuation

Nesting Population	Foraging grounds																	
	NC	EcFL	BHM	BRB	ALF	RC	FN	BA	ES	UB	AI	CB	BuA	CV	COR	ST	WA	X
Aves Island, VNZ																		
Mean	0.04	0.05	0.06	0.07	0.07	0.06	0.07	0.06	0.05	0.06	0.06	0.05	0.06	0.07	0.03	0.04	0.05	0.05
CI: 97.5%	0.13	0.19	0.20	0.20	0.21	0.20	0.21	0.21	0.18	0.17	0.20	0.16	0.19	0.25	0.09	0.15	0.15	0.18
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Buck Island																		
Mean	0.06	0.06	0.06	0.06	0.06	0.06	0.05	0.05	0.05	0.06	0.05	0.05	0.06	0.06	0.05	0.05	0.06	0.06
CI: 97.5%	0.21	0.19	0.21	0.20	0.20	0.20	0.18	0.19	0.18	0.20	0.17	0.18	0.19	0.18	0.20	0.18	0.19	0.20
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Tortuguero, CR																		
Mean	0.01	0.15	0.53	0.02	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.28
CI: 97.5%	0.04	0.33	0.84	0.04	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.64
CI: 2.5%	0.00	0.04	0.18	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Mexico																		
Mean	0.16	0.42	0.13	0.06	0.01	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.01	0.17
CI: 97.5%	0.42	0.72	0.38	0.18	0.02	0.01	0.03	0.02	0.01	0.01	0.01	0.01	0.01	0.04	0.00	0.02	0.03	0.49
CI: 2.5%	0.03	0.11	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
Southeast Cuba																		
Mean	0.27	0.11	0.11	0.07	0.03	0.03	0.03	0.03	0.01	0.02	0.01	0.01	0.02	0.05	0.01	0.03	0.04	0.12
CI: 97.5%	0.53	0.34	0.34	0.24	0.14	0.10	0.10	0.11	0.05	0.07	0.05	0.06	0.06	0.18	0.03	0.12	0.13	0.37
CI: 2.5%	0.06	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
South Florida, USA																		
Mean	0.21	0.16	0.12	0.11	0.04	0.03	0.03	0.02	0.01	0.01	0.01	0.01	0.01	0.05	0.01	0.02	0.03	0.12
CI: 97.5%	0.51	0.48	0.39	0.36	0.14	0.10	0.10	0.09	0.04	0.05	0.04	0.04	0.05	0.17	0.02	0.09	0.10	0.40
CI: 2.5%	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01
East central Florida, USA																		
Mean	0.30	0.20	0.12	0.09	0.01	0.01	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.04	0.00	0.02	0.02	0.11
CI: 97.5%	0.57	0.51	0.39	0.27	0.05	0.05	0.06	0.06	0.02	0.04	0.02	0.03	0.04	0.14	0.02	0.07	0.07	0.39
CI: 2.5%	0.04	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

698 **Table S4.** Summary of foraging ground-centric mixed stock analysis of Atlantic green turtle nesting populations (n=14) and juvenile foraging grounds (n=17).

Foraging grounds	Nesting populations													
	EcFL	SFL	MX	CR	CUB	BUC	AV	SUR	RC/N	TRI	ASC	GB	Bio	STP
North Carolina, USA														
Mean	0.19	0.10	0.35	0.16	0.08	0.00	0.01	0.02	0.00	0.01	0.02	0.05	0.01	0.00
CI: 2.5%	0.02	0.00	0.14	0.02	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
CI: 97.5%	0.41	0.26	0.59	0.36	0.18	0.00	0.05	0.06	0.01	0.03	0.06	0.11	0.02	0.01
East central Florida, USA														
Mean	0.03	0.02	0.26	0.63	0.01	0.00	0.01	0.01	0.00	0.00	0.01	0.01	0.00	0.00
CI: 2.5%	0.00	0.00	0.12	0.43	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
CI: 97.5%	0.11	0.08	0.43	0.78	0.03	0.00	0.02	0.04	0.00	0.01	0.03	0.04	0.01	0.00
Bahamas														
Mean	0.01	0.01	0.03	0.92	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.01	0.00	0.00
CI: 2.5%	0.00	0.00	0.00	0.84	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
CI: 97.5%	0.03	0.02	0.08	0.97	0.01	0.00	0.01	0.04	0.00	0.01	0.01	0.03	0.00	0.00
Barbados														
Mean	0.06	0.06	0.15	0.28	0.02	0.00	0.03	0.14	0.00	0.01	0.09	0.13	0.01	0.00
CI: 2.5%	0.00	0.00	0.02	0.08	0.00	0.00	0.00	0.03	0.00	0.00	0.01	0.01	0.00	0.00
CI: 97.5%	0.18	0.19	0.31	0.46	0.09	0.00	0.11	0.25	0.01	0.05	0.22	0.28	0.03	0.01
Almofala, Brazil														
Mean	0.01	0.03	0.02	0.14	0.02	0.00	0.04	0.19	0.00	0.02	0.26	0.25	0.01	0.00
CI: 2.5%	0.00	0.00	0.00	0.06	0.00	0.00	0.00	0.08	0.00	0.00	0.10	0.09	0.00	0.00
CI: 97.5%	0.04	0.09	0.07	0.23	0.06	0.00	0.15	0.29	0.02	0.07	0.43	0.41	0.04	0.02
Rocas Atol, Brazil														
Mean	0.02	0.03	0.02	0.05	0.02	0.00	0.06	0.21	0.01	0.05	0.37	0.16	0.02	0.01
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.05	0.00	0.00	0.15	0.01	0.00	0.00
CI: 97.5%	0.06	0.09	0.06	0.12	0.07	0.00	0.20	0.33	0.03	0.15	0.57	0.35	0.06	0.02

699

Table S4. Continuation

Foraging grounds	Nesting populations														
	EcFL	SFL	MX	CR	CUB	BUC	AV	SUR	RC/N	TRI	ASC	GB	Bio	STP	
Fernando Noronha, Brazil															
Mean	0.02	0.02	0.03	0.02	0.01	0.00	0.04	0.40	0.00	0.02	0.16	0.26	0.01	0.00	
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.26	0.00	0.00	0.02	0.09	0.00	0.00	
CI: 97.5%	0.05	0.06	0.08	0.06	0.05	0.00	0.15	0.50	0.02	0.06	0.33	0.42	0.04	0.02	
Bahia, Brazil															
Mean	0.02	0.02	0.02	0.03	0.02	0.00	0.06	0.25	0.01	0.08	0.21	0.27	0.02	0.01	
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.08	0.00	0.01	0.02	0.03	0.00	0.00	
CI: 97.5%	0.07	0.08	0.08	0.09	0.07	0.00	0.19	0.39	0.02	0.21	0.48	0.51	0.06	0.02	
Espirito Santo, Brazil															
Mean	0.01	0.01	0.01	0.01	0.01	0.00	0.05	0.26	0.01	0.04	0.29	0.29	0.02	0.01	
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.13	0.00	0.00	0.09	0.06	0.00	0.00	
CI: 97.5%	0.03	0.03	0.03	0.04	0.03	0.00	0.16	0.36	0.02	0.12	0.52	0.47	0.07	0.02	
Ubatuba, Brazil															
Mean	0.01	0.01	0.01	0.02	0.01	0.00	0.04	0.08	0.01	0.03	0.40	0.36	0.01	0.01	
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.16	0.08	0.00	0.00	
CI: 97.5%	0.04	0.04	0.04	0.05	0.04	0.00	0.13	0.17	0.02	0.10	0.68	0.60	0.05	0.02	
Arvoredo Island, Brazil															
Mean	0.01	0.01	0.01	0.01	0.01	0.00	0.06	0.16	0.01	0.07	0.37	0.27	0.01	0.01	
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.03	0.00	0.00	0.17	0.07	0.00	0.00	
CI: 97.5%	0.03	0.03	0.04	0.04	0.03	0.00	0.18	0.27	0.02	0.16	0.59	0.47	0.06	0.02	
Casino Beach, Brazil															
Mean	0.01	0.01	0.01	0.01	0.01	0.00	0.05	0.15	0.01	0.06	0.31	0.35	0.01	0.01	
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.00	0.12	0.15	0.00	0.00	
CI: 97.5%	0.03	0.04	0.04	0.04	0.03	0.00	0.16	0.26	0.03	0.15	0.53	0.55	0.06	0.03	

Foraging grounds	Nesting populations													
	EcFL	SFL	MX	CR	CUB	BUC	AV	SUR	RC/N	TRI	ASC	GB	Bio	STP
Buenos Aires, Argentina														
Mean	0.01	0.01	0.01	0.01	0.01	0.00	0.04	0.17	0.00	0.05	0.25	0.41	0.01	0.00
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.04	0.00	0.00	0.08	0.20	0.00	0.00
CI: 97.5%	0.04	0.04	0.04	0.04	0.03	0.00	0.16	0.27	0.02	0.14	0.48	0.58	0.05	0.02
Cape Verde														
Mean	0.03	0.03	0.04	0.04	0.02	0.00	0.04	0.40	0.00	0.01	0.04	0.31	0.01	0.00
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.25	0.00	0.00	0.00	0.14	0.00	0.00
CI: 97.5%	0.10	0.09	0.13	0.14	0.07	0.00	0.16	0.54	0.02	0.05	0.16	0.47	0.04	0.02
Corisco Bay, Eq. Guinea														
Mean	0.00	0.00	0.00	0.00	0.00	0.00	0.02	0.04	0.00	0.01	0.28	0.60	0.01	0.01
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.13	0.37	0.00	0.00
CI: 97.5%	0.02	0.01	0.02	0.02	0.01	0.00	0.05	0.08	0.02	0.04	0.47	0.77	0.07	0.04
Sao Tome, Sao Tome and Principe														
Mean	0.01	0.01	0.01	0.01	0.01	0.00	0.02	0.04	0.00	0.01	0.07	0.77	0.01	0.01
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.62	0.00	0.00
CI: 97.5%	0.04	0.04	0.05	0.05	0.04	0.00	0.07	0.10	0.01	0.05	0.19	0.87	0.04	0.03
West Africa: Liberia to Benin														
Mean	0.01	0.01	0.02	0.02	0.01	0.00	0.02	0.06	0.00	0.01	0.08	0.75	0.01	0.00
CI: 2.5%	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.58	0.00	0.00
CI: 97.5%	0.05	0.05	0.06	0.07	0.04	0.00	0.07	0.13	0.01	0.04	0.22	0.86	0.03	0.01




Figure S1. Comparison of mean contributions and 95% confidence intervals from Poilão rookery (West Africa) to 17 green turtle Atlantic foraging aggregations, estimated through a many-to-many mixed stock analysis, using different simulated datasets against the actual dataset - black squares. Grey circle – including a rare haplotype (Cm-A42) found at Poilão in Ascension Island sample, white triangle – including CM-A42 in Costa Rica sample, and grey diamond – adding a putative foraging ground fixed for haplotype CM-A8 (n = 99). SIM: simulated foraging ground, WA: ‘Western Africa’ – Liberia to Benin, ST: Sao Tome, COR: Corisco Bay, CV: Cape Verde, BuA: Buenos Aires, UB: Ubatuba, ALF: Almofala, CB: Cassino Beach, FN: Fernando de Noronha, ES: Espírito Santo, BA: Bahia, AI: Arvoredo Island, RC: Rocas Atol, BRB: Barbados, BHM: Bahamas, NC: North Carolina, EcFL: East central Florida. Dashed lines separate geographic regions.